

COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF
OVARIAN CANCER

CROSS REFERENCE TO RELATED APPLICATIONS

SUP A1

This application is a continuation-in-part of U.S. Application No.
5 09/617,747, filed 7/17/2000, which is a continuation-in-part of U.S. Application No.
09/404,879, filed September 24, 1999, which is a continuation-in-part of U.S. Application
No. 09/338,933, filed June 23, 1999, which is a continuation-in-part of U.S. Application
Nos. 09/216,003, filed December 17, 1999, and 09/215,681, filed December 17, 1998.

TECHNICAL FIELD

10 The present invention relates generally to ovarian cancer therapy. The
invention is more specifically related to polypeptides comprising at least a portion of an
ovarian carcinoma protein, and to polynucleotides encoding such polypeptides, as well as
antibodies and immune system cells that specifically recognize such polypeptides. Such
polypeptides, polynucleotides, antibodies and cells may be used in vaccines and
15 pharmaceutical compositions for treatment of ovarian cancer.

BACKGROUND OF THE INVENTION

Ovarian cancer is a significant health problem for women in the United
States and throughout the world. Although advances have been made in detection and
therapy of this cancer, no vaccine or other universally successful method for prevention or
20 treatment is currently available. Management of the disease currently relies on a
combination of early diagnosis and aggressive treatment, which may include one or more
of a variety of treatments such as surgery, radiotherapy, chemotherapy and hormone
therapy. The course of treatment for a particular cancer is often selected based on a variety
of prognostic parameters, including an analysis of specific tumor markers. However, the
25 use of established markers often leads to a result that is difficult to interpret, and high
mortality continues to be observed in many cancer patients.

09636301.081000

Immunotherapies have the potential to substantially improve cancer treatment and survival. Such therapies may involve the generation or enhancement of an immune response to an ovarian carcinoma antigen. However, to date, relatively few ovarian carcinoma antigens are known and the generation of an immune response against such antigens has not been shown to be therapeutically beneficial.

Accordingly, there is a need in the art for improved methods for identifying ovarian tumor antigens and for using such antigens in the therapy of ovarian cancer. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

Briefly stated, this invention provides compositions and methods for the therapy of cancer, such as ovarian cancer. In one aspect, the present invention provides polypeptides comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished. Within certain embodiments, the ovarian carcinoma protein comprises a sequence that is encoded by a polynucleotide sequence selected from the group consisting of SEQ ID NOs:1-81, 313-331, 359, 366, 379, 385-387, 391 and complements of such polynucleotides.

The present invention further provides polynucleotides that encode a polypeptide as described above or a portion thereof, expression vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors.

Within other aspects, the present invention provides pharmaceutical compositions and vaccines. Pharmaceutical compositions may comprise a physiologically acceptable carrier or excipient in combination with one or more of: (i) a polypeptide comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid

sequence encoded by a polynucleotide that comprises a sequence recited in any one of SEQ ID NOs:1-81, 313-331, 359, 366, 379, 385-387 or 391; (ii) a polynucleotide encoding such a polypeptide; (iii) an antibody that specifically binds to such a polypeptide; (iv) an antigen-presenting cell that expresses such a polypeptide and/or (v) a T cell that specifically reacts with such a polypeptide. Vaccines may comprise a non-specific immune response enhancer in combination with one or more of: (i) a polypeptide comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid sequence encoded by a polynucleotide that comprises a sequence recited in any one of SEQ ID NOs:1-81, 313-331, 359, 366, 379, 385-387 or 391; (ii) a polynucleotide encoding such a polypeptide; (iii) an anti-idiotypic antibody that is specifically bound by an antibody that specifically binds to such a polypeptide; (iv) an antigen-presenting cell that expresses such a polypeptide and/or (v) a T cell that specifically reacts with such a polypeptide.

The present invention further provides, in other aspects, fusion proteins that comprise at least one polypeptide as described above, as well as polynucleotides encoding such fusion proteins.

Within related aspects, pharmaceutical compositions comprising a fusion protein or polynucleotide encoding a fusion protein in combination with a physiologically acceptable carrier are provided.

Vaccines are further provided, within other aspects, comprising a fusion protein or polynucleotide encoding a fusion protein in combination with a non-specific immune response enhancer.

Within further aspects, the present invention provides methods for inhibiting the development of a cancer in a patient, comprising administering to a patient a pharmaceutical composition or vaccine as recited above.

The present invention further provides, within other aspects, methods for stimulating and/or expanding T cells, comprising contacting T cells with (a) a polypeptide

comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid sequence encoded by a polynucleotide that comprises a sequence recited in any one of SEQ ID NOs:1-387 or 391; (b) a polynucleotide encoding such a polypeptide and/or (c) an antigen presenting cell that expresses such a polypeptide under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells. Such polypeptide, polynucleotide and/or antigen presenting cell(s) may be present within a pharmaceutical composition or vaccine, for use in stimulating and/or expanding T cells in a mammal.

Within other aspects, the present invention provides methods for inhibiting the development of ovarian cancer in a patient, comprising administering to a patient T cells prepared as described above.

Within further aspects, the present invention provides methods for inhibiting the development of ovarian cancer in a patient, comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a polypeptide comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid sequence encoded by a polynucleotide that comprises a sequence recited in any one of SEQ ID NOs: 1-387 or 391; (ii) a polynucleotide encoding such a polypeptide; or (iii) an antigen-presenting cell that expresses such a polypeptide; such that T cells proliferate; and (b) administering to the patient an effective amount of the proliferated T cells, and thereby inhibiting the development of ovarian cancer in the patient. The proliferated cells may be cloned prior to administration to the patient.

The present invention also provides, within other aspects, methods for identifying secreted tumor antigens. Such methods comprise the steps of: (a) implanting tumor cells in an immunodeficient mammal; (b) obtaining serum from the immunodeficient

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

Figure 3 presents results of microarray expression analysis of the ovarian carcinoma sequence designated O8E.

Figure 5 presents the ovarian carcinoma polynucleotide designated 3f (SEQ ID NO:76).

Figures 7A and 7B present the ovarian carcinoma polynucleotides designated 8e (SEQ ID NO:78) and 8h (SEQ ID NO:79).

Figure 9 presents the ovarian carcinoma polynucleotide designated 12h (SEQ ID NO:81).

Figure 11 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 6b.

Figure 13 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 12c.

Figure 14 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 12h.

Figures 15A-15EEE depict partial sequences of additional polynucleotides
25 encoding representative secreted ovarian carcinoma antigens (SEQ ID NOs:82-310).

Figure 16 is a diagram illustrating the location of various partial O8E sequences within the full length sequence.

Figure 17 is a graph illustrating the results of epitope mapping studies on O8E protein.

Figure 18 is graph of a fluorescence activated cell sorting (FACS) analysis of O8E cell surface expression.

Figure 19 is graph of a FACS analysis of O8E cell surface expression.

Figure 20 shows FACS analysis results for O8E transfected HEK293 cells demonstrating cell surface expression of O8E.

Figure 21 shows FACS analysis results for SKBR3 breast tumor cells demonstrating cell surface expression of O8E.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the therapy of cancer, such as ovarian cancer. The compositions described herein may include immunogenic polypeptides, polynucleotides encoding such polypeptides, binding agents such as antibodies that bind to a polypeptide, antigen presenting cells (APCs) and/or immune system cells (*e.g.*, T cells).

Polypeptides of the present invention generally comprise at least an immunogenic portion of an ovarian carcinoma protein or a variant thereof. Certain ovarian carcinoma proteins have been identified using an immunoassay technique, and are referred to herein as ovarian carcinoma antigens. An "ovarian carcinoma antigen" is a protein that is expressed by ovarian tumor cells (preferably human cells) at a level that is at least two fold higher than the level in normal ovarian cells. Certain ovarian carcinoma antigens react detectably (within an immunoassay, such as an ELISA or Western blot) with antisera generated against serum from an immunodeficient animal implanted with a human ovarian tumor. Such ovarian carcinoma antigens are shed or secreted from an ovarian tumor into the sera of the immunodeficient animal. Accordingly, certain ovarian carcinoma antigens provided herein are secreted antigens. Certain nucleic acid sequences of the subject invention generally comprise a DNA or RNA sequence that encodes all or a portion of such a polypeptide, or that is complementary to such a sequence.

The present invention further provides ovarian carcinoma sequences that are identified using techniques to evaluate altered expression within an ovarian tumor. Such sequences may be polynucleotide or protein sequences. Ovarian carcinoma sequences are generally expressed in an ovarian tumor at a level that is at least two fold, and preferably at least five fold, greater than the level of expression in normal ovarian tissue, as determined using a representative assay provided herein. Certain partial ovarian carcinoma polynucleotide sequences are presented herein. Proteins encoded by genes comprising such polynucleotide sequences (or complements thereof) are also considered ovarian carcinoma proteins.

Antibodies are generally immune system proteins, or antigen-binding fragments thereof, that are capable of binding to at least a portion of an ovarian carcinoma polypeptide as described herein. T cells that may be employed within the compositions provided herein are generally T cells (*e.g.*, CD4⁺ and/or CD8⁺) that are specific for such a polypeptide. Certain methods described herein further employ antigen-presenting cells (such as dendritic cells or macrophages) that express an ovarian carcinoma polypeptide as provided herein.

OVARIAN CARCINOMA POLYNUCLEOTIDES

Any polynucleotide that encodes an ovarian carcinoma protein or a portion or other variant thereof as described herein is encompassed by the present invention. Preferred polynucleotides comprise at least 15 consecutive nucleotides, preferably at least 30 consecutive nucleotides, and more preferably at least 45 consecutive nucleotides, that encode a portion of an ovarian carcinoma protein. More preferably, a polynucleotide encodes an immunogenic portion of an ovarian carcinoma protein, such as an ovarian carcinoma antigen. Polynucleotides complementary to any such sequences are also encompassed by the present invention. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. Additional coding or non-coding sequences may, but need not, be present

Polynucleotides may comprise a native sequence (*i.e.*, an endogenous sequence that encodes an ovarian carcinoma protein or a portion thereof) or may comprise a variant of such a sequence. Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions such that the immunogenicity of the encoded polypeptide is not diminished, relative to a native ovarian carcinoma protein. The effect on the immunogenicity of the encoded polypeptide may generally be assessed as described herein. Variants preferably exhibit at least about 70% identity, more preferably at least about 80% identity and most preferably at least about 90% identity to a polynucleotide sequence that encodes a native ovarian carcinoma protein or a portion thereof.

15 those of ordinary skill in the art, such as Megalign, using default parameters. Comparisons
between two sequences are typically performed by comparing the sequences over a
comparison window to identify and compare local regions of sequence similarity. A
“comparison window” as used herein, refers to a segment of at least about 20 contiguous
positions, usually 30 to about 75, or 40 to about 50, in which a sequence may be compared
20 to a reference sequence of the same number of contiguous positions after the two sequences
are optimally aligned. Optimal alignment of sequences for comparison may be conducted,
for example, using the Megalign program in the Lasergene suite of bioinformatics software
(DNASTAR, Inc., Madison, WI), using default parameters. Preferably, the percentage of
sequence identity is determined by comparing two optimally aligned sequences over a
25 window of comparison of at least 20 positions, wherein the portion of the polynucleotide or
polypeptide sequence in the window may comprise additions or deletions (*i.e.*, gaps) of 20
% or less, usually 5 to 15 %, or 10 to 12%, relative to the reference sequence (which does
not contain additions or deletions). The percent identity may be calculated by determining
the number of positions at which the identical nucleic acid bases or amino acid residue

occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

5 Variants may also, or alternatively, be substantially homologous to a native gene, or a portion or complement thereof. Such polynucleotide variants are capable of hybridizing under moderately stringent conditions to a naturally occurring DNA sequence encoding a native ovarian carcinoma protein (or a complementary sequence). Suitable moderately stringent conditions include prewashing in a solution of 5 X SSC, 0.5% SDS,
10 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS.

It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a
15 polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of
20 one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

Polynucleotides may be prepared using any of a variety of techniques. For
25 example, an ovarian carcinoma polynucleotide may be identified, as described in more detail below, by screening a late passage ovarian tumor expression library with antisera generated against sera of immunocompetent mice after injection of such mice with sera from SCID mice implanted with late passage ovarian tumors. Ovarian carcinoma polynucleotides may also be identified using any of a variety of techniques designed to

5 An amplified portion may be used to isolate a full length gene from a suitable library (*e.g.*, an ovarian carcinoma cDNA library) using well known techniques. Within such techniques, a library (cDNA or genomic) is screened using one or more polynucleotide probes or primers suitable for amplification. Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for
10 identifying 5' and upstream regions of genes. Genomic libraries are preferred for obtaining introns and extending 5' sequences.

For hybridization techniques, a partial sequence may be labeled (e.g., by nick-translation or end-labeling with ^{32}P) using well known techniques. A bacterial or bacteriophage library is then screened by hybridizing filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe (see Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989). Hybridizing colonies or plaques are selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and partial sequences may be generated to identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may involve generating a series of deletion clones. The resulting overlapping sequences are then assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

11

One such amplification technique is inverse PCR (*see* Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularized by intramolecular ligation and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Additional techniques include capture PCR (Lagerstrom et al., *PCR Methods Applic.* 1:111-19, 1991) and walking PCR (Parker et al., *Nucl. Acids. Res.* 19:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

In certain instances, it is possible to obtain a full length cDNA sequence by analysis of sequences provided in an expressed sequence tag (EST) database, such as that available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (*e.g.*, NCBI BLAST searches), and such ESTs may be used to generate a contiguous full length sequence.

25

5

10

15

25

Real-time PCR may be performed, for example, using a Perkin Elmer/Applied Biosystems (Foster City, CA) 7700 Prism instrument. Matching primers and fluorescent probes may be designed for genes of interest using, for example, the primer express program provided by Perkin Elmer/Applied Biosystems (Foster City, CA). Optimal concentrations of primers and probes may be initially determined by those of ordinary skill in the art, and control (e.g., β -actin) primers and probes may be obtained commercially from, for example, Perkin Elmer/Applied Biosystems (Foster City, CA). To quantitate the amount of specific RNA in a sample, a standard curve is generated alongside using a plasmid containing the gene of interest. Standard curves may be generated using the Ct values determined in the real-time PCR, which are related to the initial cDNA concentration used in the assay. Standard dilutions ranging from 10^{-10} to 10^{-6} copies of the gene of interest are generally sufficient. In addition, a standard curve is generated for the control sequence. This permits standardization of initial RNA content of a tissue sample to the amount of control for comparison purposes.

Polynucleotide variants may generally be prepared by any method known in the art, including chemical synthesis by, for example, solid phase phosphoramidite chemical synthesis. Modifications in a polynucleotide sequence may also be introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis (*see* Adelman et al., *DNA* 2:183, 1983). Alternatively, RNA molecules may be generated by *in vitro* or *in vivo* transcription of DNA sequences encoding an ovarian carcinoma antigen, or portion thereof, provided that the DNA is incorporated into a vector with a suitable RNA polymerase promoter (such as T7 or SP6). Certain portions may be used to prepare an encoded polypeptide, as described herein. In addition, or alternatively, a portion may be administered to a patient such that the encoded polypeptide is generated *in vivo*.

A portion of a sequence complementary to a coding sequence (*i.e.*, an antisense polynucleotide) may also be used as a probe or to modulate gene expression. cDNA constructs that can be transcribed into antisense RNA may also be introduced into cells or tissues to facilitate the production of antisense RNA. An antisense polynucleotide

may be used, as described herein, to inhibit expression of an ovarian carcinoma protein. Antisense technology can be used to control gene expression through triple-helix formation, which compromises the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors or regulatory molecules (*see* Gee et al., *In* 5 Huber and Carr, *Molecular and Immunologic Approaches*, Futura Publishing Co. (Mt. Kisco, NY; 1994). Alternatively, an antisense molecule may be designed to hybridize with a control region of a gene (*e.g.*, promoter, enhancer or transcription initiation site), and block transcription of the gene; or to block translation by inhibiting binding of a transcript to ribosomes.

10 Any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl- methyl-, thio- and other 15 modified forms of adenine, cytidine, guanine, thymine and uridine.

Nucleotide sequences as described herein may be joined to a variety of other nucleotide sequences using established recombinant DNA techniques. For example, a polynucleotide may be cloned into any of a variety of cloning vectors, including plasmids, phagemids, lambda phage derivatives and cosmids. Vectors of particular interest include 20 expression vectors, replication vectors, probe generation vectors and sequencing vectors. In general, a vector will contain an origin of replication functional in at least one organism, convenient restriction endonuclease sites and one or more selectable markers. Other elements will depend upon the desired use, and will be apparent to those of ordinary skill in the art.

25 Within certain embodiments, polynucleotides may be formulated so as to permit entry into a cell of a mammal, and expression therein. Such formulations are particularly useful for therapeutic purposes, as described below. Those of ordinary skill in the art will appreciate that there are many ways to achieve expression of a polynucleotide in a target cell, and any suitable method may be employed. For example, a polynucleotide

may be incorporated into a viral vector such as, but not limited to, adenovirus, adeno-associated virus, retrovirus, or vaccinia or other pox virus (e.g., avian pox virus). Techniques for incorporating DNA into such vectors are well known to those of ordinary skill in the art. A retroviral vector may additionally transfer or incorporate a gene for a selectable marker (to aid in the identification or selection of transduced cells) and/or a targeting moiety, such as a gene that encodes a ligand for a receptor on a specific target cell, to render the vector target specific. Targeting may also be accomplished using an antibody, by methods known to those of ordinary skill in the art.

Other formulations for therapeutic purposes include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. A preferred colloidal system for use as a delivery vehicle *in vitro* and *in vivo* is a liposome (i.e., an artificial membrane vesicle). The preparation and use of such systems is well known in the art.

OVARIAN CARCINOMA POLYPEPTIDES

Within the context of the present invention, polypeptides may comprise at least an immunogenic portion of an ovarian carcinoma protein or a variant thereof, as described herein. As noted above, certain ovarian carcinoma proteins are ovarian carcinoma antigens that are expressed by ovarian tumor cells and react detectably within an immunoassay (such as an ELISA) with antisera generated against serum from an immunodeficient animal implanted with an ovarian tumor. Other ovarian carcinoma proteins are encoded by ovarian carcinoma polynucleotides recited herein. Polypeptides as described herein may be of any length. Additional sequences derived from the native protein and/or heterologous sequences may be present, and such sequences may (but need not) possess further immunogenic or antigenic properties.

An "immunogenic portion," as used herein is a portion of an antigen that is recognized (i.e., specifically bound) by a B-cell and/or T-cell surface antigen receptor. Such immunogenic portions generally comprise at least 5 amino acid residues, more

preferably at least 10, and still more preferably at least 20 amino acid residues of an ovarian carcinoma protein or a variant thereof. Preferred immunogenic portions are encoded by cDNA molecules isolated as described herein. Further immunogenic portions may generally be identified using well known techniques, such as those summarized in Paul, 5 *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 1993) and references cited therein. Such techniques include screening polypeptides for the ability to react with ovarian carcinoma protein-specific antibodies, antisera and/or T-cell lines or clones. As used herein, antisera and antibodies are "ovarian carcinoma protein-specific" if they specifically bind to an ovarian carcinoma protein (*i.e.*, they react with the ovarian carcinoma protein in an ELISA or other immunoassay, and do not react detectably with 10 unrelated proteins). Such antisera, antibodies and T cells may be prepared as described herein, and using well known techniques. An immunogenic portion of a native ovarian carcinoma protein is a portion that reacts with such antisera, antibodies and/or T-cells at a level that is not substantially less than the reactivity of the full length polypeptide (*e.g.*, in 15 an ELISA and/or T-cell reactivity assay). Such immunogenic portions may react within such assays at a level that is similar to or greater than the reactivity of the full length protein. Such screens may generally be performed using methods well known to those of ordinary skill in the art, such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. For example, a polypeptide 20 may be immobilized on a solid support and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A.

As noted above, a composition may comprise a variant of a native ovarian carcinoma protein. A polypeptide "variant," as used herein, is a polypeptide that differs 25 from a native ovarian carcinoma protein in one or more substitutions, deletions, additions and/or insertions, such that the immunogenicity of the polypeptide is not substantially diminished. In other words, the ability of a variant to react with ovarian carcinoma protein-specific antisera may be enhanced or unchanged, relative to the native ovarian carcinoma protein, or may be diminished by less than 50%, and preferably less than 20%, relative to

5

10

15

20

25

As noted above, polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs

Polypeptides may be prepared using any of a variety of well known techniques. Recombinant polypeptides encoded by DNA sequences as described above may be readily prepared from the DNA sequences using any of a variety of expression vectors known to those of ordinary skill in the art. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line such as COS or CHO. Supernatants from suitable host/vector systems which secrete recombinant protein or polypeptide into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant polypeptide.

Within certain specific embodiments, a polypeptide may be a fusion protein that comprises multiple polypeptides as described herein, or that comprises one polypeptide as described herein and a known tumor antigen, such as an ovarian carcinoma protein or a

variant of such a protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both
 5 immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the protein.

Fusion proteins may generally be prepared using standard techniques,
 10 including chemical conjugation. Preferably, a fusion protein is expressed as a recombinant protein, allowing the production of increased levels, relative to a non-fused protein, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide
 15 linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion protein that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptide components by a distance sufficient to ensure that each polypeptide
 20 folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides;
 25 and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA*

5 interference.

10 present 3' to the DNA sequence encoding the second polypeptide.

15 336:86-91, 1997).

may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen present cells. Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known

as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at the amino terminus has been described (see *Biotechnology* 10:795-798, 1992). Within a preferred embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

In general, polypeptides (including fusion proteins) and polynucleotides as described herein are isolated. An "isolated" polypeptide or polynucleotide is one that is removed from its original environment. For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

20 BINDING AGENTS

The present invention further provides agents, such as antibodies and antigen-binding fragments thereof, that specifically bind to an ovarian carcinoma protein. As used herein, an antibody, or antigen-binding fragment thereof, is said to "specifically bind" to an ovarian carcinoma protein if it reacts at a detectable level (within, for example, an ELISA) with an ovarian carcinoma protein, and does not react detectably with unrelated proteins under similar conditions. As used herein, "binding" refers to a noncovalent association between two separate molecules such that a "complex" is formed. The ability to bind may be evaluated by, for example, determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the

complex is divided by the product of the component concentrations. In general, two compounds are said to "bind," in the context of the present invention, when the binding constant for complex formation exceeds about 10^3 L/mol. The binding constant may be determined using methods well known in the art.

5 Binding agents may be further capable of differentiating between patients with and without a cancer, such as ovarian cancer, using the representative assays provided herein. In other words, antibodies or other binding agents that bind to a ovarian carcinoma antigen will generate a signal indicating the presence of a cancer in at least about 20% of patients with the disease, and will generate a negative signal indicating the absence of the
10 disease in at least about 90% of individuals without the cancer. To determine whether a binding agent satisfies this requirement, biological samples (*e.g.*, blood, sera, leukophoresis, urine and/or tumor biopsies) from patients with and without a cancer (as determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. It will be apparent that a
15 statistically significant number of samples with and without the disease should be assayed. Each binding agent should satisfy the above criteria; however, those of ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component, an
20 RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation of
25 monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen without

5

10

25

extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Within certain embodiments, the use of antigen-binding fragments of antibodies may be preferred. Such fragments include Fab fragments, which may be prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by affinity chromatography on protein A bead columns.

Monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, Shigella toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (e.g., covalently bonded) to a suitable monoclonal antibody either directly or indirectly (e.g., via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (e.g., a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also

facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also

bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

Also provided herein are anti-idiotypic antibodies that mimic an immunogenic portion of an ovarian carcinoma protein. Such antibodies may be raised against an antibody, or antigen-binding fragment thereof, that specifically binds to an immunogenic portion of an ovarian carcinoma protein, using well known techniques. Anti-idiotypic antibodies that mimic an immunogenic portion of an ovarian carcinoma protein are those antibodies that bind to an antibody, or antigen-binding fragment thereof, that specifically binds to an immunogenic portion of an ovarian carcinoma protein, as described herein.

T CELLS

Immunotherapeutic compositions may also, or alternatively, comprise T cells specific for an ovarian carcinoma protein. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example, T cells may be present within (or isolated from) bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood of a mammal, such as a patient, using a commercially available cell separation system, such as the CEPRATE™ system, available from CellPro Inc., Bothell WA (see also

U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or unrelated humans, non-human animals, cell lines or cultures.

T cells may be stimulated with an ovarian carcinoma polypeptide, polynucleotide encoding an ovarian carcinoma polypeptide and/or an antigen presenting cell (APC) that expresses such a polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific for the polypeptide. Preferably, an ovarian carcinoma polypeptide or polynucleotide is present within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

T cells are considered to be specific for an ovarian carcinoma polypeptide if the T cells kill target cells coated with an ovarian carcinoma polypeptide or expressing a gene encoding such a polypeptide. T cell specificity may be evaluated using any of a variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen et al., *Cancer Res.* 54:1065-1070, 1994. Alternatively, detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (e.g., by pulse-labeling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with an ovarian carcinoma polypeptide (200 ng/ml - 100 µg/ml, preferably 100 ng/ml - 25 µg/ml) for 3 - 7 days should result in at least a two fold increase in proliferation of the T cells and/or contact as described above for 2-3 hours should result in activation of the T cells, as measured using standard cytokine assays in which a two fold increase in the level of cytokine release (e.g., TNF or IFN-γ) is indicative of T cell activation (see Coligan et al., *Current Protocols in Immunology*, vol. 1, Wiley Interscience (Greene 1998). T cells that have been activated in response to an ovarian carcinoma polypeptide, polynucleotide or ovarian carcinoma polypeptide-expressing APC may be CD4⁺ and/or CD8⁺. Ovarian

carcinoma polypeptide-specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient or a related or unrelated donor and are administered to the patient following stimulation and expansion.

For therapeutic purposes, CD4⁺ or CD8⁺ T cells that proliferate in response to an ovarian carcinoma polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to an ovarian carcinoma polypeptide, with or without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesize an ovarian carcinoma polypeptide. Alternatively, one or more T cells that proliferate in the presence of an ovarian carcinoma polypeptide can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting dilution. Following expansion, the cells may be administered back to the patient as described, for example, by Chang et al., *Crit. Rev. Oncol. Hematol.* 22:213, 1996.

15

PHARMACEUTICAL COMPOSITIONS AND VACCINES

Within certain aspects, polypeptides, polynucleotides, binding agents and/or immune system cells as described herein may be incorporated into pharmaceutical compositions or vaccines. Pharmaceutical compositions comprise one or more such compounds or cells and a physiologically acceptable carrier. Vaccines may comprise one or more such compounds or cells and a non-specific immune response enhancer. A non-specific immune response enhancer may be any substance that enhances an immune response to an exogenous antigen. Examples of non-specific immune response enhancers include adjuvants, biodegradable microspheres (*e.g.*, polylactic galactide) and liposomes (into which the compound is incorporated; *see e.g.*, Fullerton, U.S. Patent No. 4,235,877). Vaccine preparation is generally described in, for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995). Pharmaceutical compositions and vaccines within the scope of the present invention may also contain other compounds, which may be biologically active or inactive. For example,

A pharmaceutical composition or vaccine may contain DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in*

situ. As noted above, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface. In a preferred embodiment, the DNA may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *PNAS* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *PNAS* 91:215-219, 1994; Kass-Eisler et al., *PNAS* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. Compositions of the present invention may be

formulated for any appropriate manner of administration, including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral
 5 administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S.
 10 Patent Nos. 4,897,268 and 5,075,109.

Such compositions may also comprise buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide) and/or
 15 preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

Any of a variety of non-specific immune response enhancers may be employed in the vaccines of this invention. For example, an adjuvant may be included.
 20 Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI), Merck Adjuvant 65 (Merck and
 25 Company, Inc., Rahway, NJ), alum, biodegradable microspheres, monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF or interleukin-2, -7, or -12, may also be used as adjuvants.

Within the vaccines provided herein, the adjuvant composition is preferably designed to induce an immune response predominantly of the Th1 type. High levels of

Th1-type cytokines (e.g., IFN- γ , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (e.g., IL-4, IL-5, IL-6, IL-10 and TNF- β) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL), together with an aluminum salt. MPL adjuvants are available from Ribi ImmunoChem Research Inc. (Hamilton, MT; see US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). Also preferred is AS-2 (SmithKline Beecham). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555. Another preferred adjuvant is a saponin, preferably QS21, which may be used alone or in combination with other adjuvants. For example, an enhanced system involves the combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprises an oil-in-water emulsion and tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210. Any vaccine provided herein may be prepared using well known methods that result in a combination of antigen, immune response enhancer and a suitable carrier or excipient.

The compositions described herein may be administered as part of a sustained release formulation (*i.e.*, a formulation such as a capsule or sponge that effects a slow release of compound following administration). Such formulations may generally be prepared using well known technology and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain a polypeptide, polynucleotide or antibody dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane. Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

Any of a variety of delivery vehicles may be employed within pharmaceutical compositions and vaccines to facilitate production of an antigen-specific immune response that targets tumor cells. Delivery vehicles include antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response, to have anti-tumor effects *per se* and/or to be immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (*see* Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*, with marked cytoplasmic processes (dendrites) visible *in vitro*) and based on the lack of differentiation

5

10

20

25

place *ex vivo*, and a composition or vaccine comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic

5 cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant

10 bacterium or viruses (*e.g.*, vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (*e.g.*, a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

15

CANCER THERAPY

In further aspects of the present invention, the compositions described herein may be used for immunotherapy of cancer, such as ovarian cancer. Within such methods, pharmaceutical compositions and vaccines are typically administered to a patient.

20 As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may or may not be afflicted with cancer. Accordingly, the above pharmaceutical compositions and vaccines may be used to prevent the development of a cancer or to treat a patient afflicted with a cancer. Within certain preferred embodiments, a patient is afflicted with ovarian cancer. Such cancer may be diagnosed using criteria generally accepted in the

25 art, including the presence of a malignant tumor. Pharmaceutical compositions and vaccines may be administered either prior to or following surgical removal of primary tumors and/or treatment such as administration of radiotherapy or conventional chemotherapeutic drugs.

5 Within other embodiments, immunotherapy may be passive immunotherapy, in which treatment involves the delivery of agents with established tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T lymphocytes (such as CD8⁺ cytotoxic T lymphocytes and CD4⁺ T-helper tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells and lymphokine-activated killer cells), B cells and antigen-presenting cells (such as dendritic cells and macrophages) expressing a polypeptide provided herein. T cell receptors and antibody receptors specific for the polypeptides recited herein may be cloned, expressed and transferred into other vectors or effector cells for adoptive immunotherapy. The polypeptides provided herein may also be used to generate antibodies or anti-idiotypic antibodies (as described above and in U.S. Patent No. 4,918,164) for passive immunotherapy.

36

system. Cultured effector cells for use in therapy must be able to grow and distribute widely, and to survive long term *in vivo*. Studies have shown that cultured effector cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (*see*, for example, Cheever et al.,
5 *Immunological Reviews* 157:177, 1997).

Alternatively, a vector expressing a polypeptide recited herein may be introduced into stem cells taken from a patient and clonally propagated *in vitro* for autologous transplant back into the same patient.

Routes and frequency of administration, as well as dosage, will vary from
10 individual to individual, and may be readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration), orally or in the bed of a resected tumor. Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at
15 intervals of 1 month, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response, and is at least 10-50% above the basal (*i.e.*, untreated) level.. Such response can be monitored by measuring the anti-tumor antibodies
20 in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing the patient's tumor cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (*e.g.*, more frequent remissions, complete or partial or longer disease-free survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and
25 vaccines comprising one or more polypeptides, the amount of each polypeptide present in a dose ranges from about 100 µg to 5 mg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit.

Such a response can be monitored by establishing an improved clinical outcome (e.g., more frequent remissions, complete or partial, or longer disease-free survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to an ovarian carcinoma antigen generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

SCREENS FOR IDENTIFYING SECRETED OVARIAN CARCINOMA ANTIGENS

The present invention provides methods for identifying secreted tumor antigens. Within such methods, tumors are implanted into immunodeficient animals such as SCID mice and maintained for a time sufficient to permit secretion of tumor antigens into serum. In general, tumors may be implanted subcutaneously or within the gonadal fat pad of an immunodeficient animal and maintained for 1-9 months, preferably 1-4 months. Implantation may generally be performed as described in WO 97/18300. The serum containing secreted antigens is then used to prepare antisera in immunocompetent mice, using standard techniques and as described herein. Briefly, 50-100 μ L of sera (pooled from three sets of immunodeficient mice, each set bearing a different SCID-derived human ovarian tumor) may be mixed 1:1 (vol:vol) with an appropriate adjuvant, such as RIBI-MPL or MPL + TDM (Sigma Chemical Co., St. Louis, MO) and injected intraperitoneally into syngeneic immunocompetent animals at monthly intervals for a total of 5 months. Antisera from animals immunized in such a manner may be obtained by drawing blood after the third, fourth and fifth immunizations. The resulting antiserum is generally pre-cleared of *E. coli* and phage antigens and used (generally following dilution, such as 1:200) in a serological expression screen.

The library is typically an expression library containing cDNAs from one or more tumors of the type that was implanted into SCID mice. This expression library may be prepared in any suitable vector, such as λ -screen (Novagen). cDNAs that encode a polypeptide that reacts with the antiserum may be identified using standard techniques, and

sequenced. Such cDNA molecules may be further characterized to evaluate expression in tumor and normal tissue, and to evaluate antigen secretion in patients.

The methods provided herein have advantages over other methods for tumor antigen discovery. In particular, all antigens identified by such methods should be secreted or released through necrosis of the tumor cells. Such antigens may be present on the surface of tumor cells for an amount of time sufficient to permit targeting and killing by the immune system, following vaccination.

METHODS FOR DETECTING CANCER

In general, a cancer may be detected in a patient based on the presence of one or more ovarian carcinoma proteins and/or polynucleotides encoding such proteins in a biological sample (such as blood, sera, urine and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a cancer such as ovarian cancer. In addition, such proteins may be useful for the detection of other cancers. The binding agents provided herein generally permit detection of the level of protein that binds to the agent in the biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding a tumor protein, which is also indicative of the presence or absence of a cancer. In general, an ovarian carcinoma-associated sequence should be present at a level that is at least three fold higher in tumor tissue than in normal tissue

There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample. *See, e.g.,* Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, the presence or absence of a cancer in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b) detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

In a preferred embodiment, the assay involves the use of binding agent immobilized on a solid support to bind to and remove the polypeptide from the remainder

of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include full length ovarian carcinoma proteins and portions thereof to which the binding agent binds, as described above.

The solid support may be any material known to those of ordinary skill in the art to which the tumor protein may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about

10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both
5 the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

10 In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a
15 detection reagent (preferably a second antibody capable of binding to a different site on the polypeptide) containing a reporter group is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as
20 described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered
25 saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.,* incubation time) is a period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with ovarian cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art

will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include those groups recited above.

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of a cancer, such as ovarian cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value for the detection of a cancer is the average mean signal obtained when the immobilized antibody is incubated with samples from patients without the cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for the cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this

embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for a cancer.

10 In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the binding agent is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized binding agent as the sample passes through the membrane. A second, labeled binding agent then binds to the binding agent-polypeptide complex as a solution containing
15 the second binding agent flows through the membrane. The detection of bound second binding agent may then be performed as described above. In the strip test format, one end of the membrane to which binding agent is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second binding agent and to the area of immobilized binding agent. Concentration of second
20 binding agent at the area of immobilized antibody indicates the presence of a cancer. Typically, the concentration of second binding agent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of binding agent immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of
25 polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferred binding agents for use in such assays are antibodies and antigen-binding fragments thereof. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more

preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the tumor proteins or binding agents of the present invention. The above descriptions are intended to be exemplary only. For example, it will be apparent to those of ordinary skill in the art that the above protocols may be readily modified to use ovarian carcinoma polypeptides to detect antibodies that bind to such polypeptides in a biological sample. The detection of such ovarian carcinoma protein specific antibodies may correlate with the presence of a cancer.

A cancer may also, or alternatively, be detected based on the presence of T cells that specifically react with an ovarian carcinoma protein in a biological sample. Within certain methods, a biological sample comprising CD4⁺ and/or CD8⁺ T cells isolated from a patient is incubated with an ovarian carcinoma protein, a polynucleotide encoding such a polypeptide and/or an APC that expresses at least an immunogenic portion of such a polypeptide, and the presence or absence of specific activation of the T cells is detected. Suitable biological samples include, but are not limited to, isolated T cells. For example, T cells may be isolated from a patient by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with an ovarian carcinoma protein (*e.g.*, 5 - 25 µg/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence of ovarian carcinoma protein to serve as a control. For CD4⁺ T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8⁺ T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a cancer in the patient.

As noted above, a cancer may also, or alternatively, be detected based on the level of mRNA encoding an ovarian carcinoma protein in a biological sample. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of an ovarian carcinoma protein cDNA

derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for (*i.e.*, hybridizes to) a polynucleotide encoding the ovarian carcinoma protein. The amplified cDNA is then separated and detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes that specifically hybridize to a polynucleotide encoding an ovarian carcinoma protein may be used in a hybridization assay to detect the presence of polynucleotide encoding the tumor protein in a biological sample.

To permit hybridization under assay conditions, oligonucleotide primers and probes should comprise an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding an ovarian carcinoma protein that is at least 10 nucleotides, and preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers and/or probes hybridize to a polynucleotide encoding a polypeptide described herein under moderately stringent conditions, as defined above. Oligonucleotide primers and/or probes which may be usefully employed in the diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule having a sequence provided herein. Techniques for both PCR based assays and hybridization assays are well known in the art (*see*, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989).

One preferred assay employs RT-PCR, in which PCR is applied in conjunction with reverse transcription. Typically, RNA is extracted from a biological sample such as a biopsy tissue and is reverse transcribed to produce cDNA molecules. PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a test patient and from an individual who is not afflicted with a cancer. The amplification reaction may be performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in

In another embodiment, ovarian carcinoma proteins and polynucleotides encoding such proteins may be used as markers for monitoring the progression of cancer.

Certain *in vivo* diagnostic assays may be performed directly on a tumor. One such assay involves contacting tumor cells with a binding agent. The bound binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

25 DIAGNOSTIC KITS

46

antibody or fragment thereof that specifically binds to an ovarian carcinoma protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection
5 reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

Alternatively, a kit may be designed to detect the level of mRNA encoding an ovarian carcinoma protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer, as described above, that hybridizes to a
10 polynucleotide encoding an ovarian carcinoma protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding an ovarian carcinoma protein.

15 The following Examples are offered by way of illustration and not by way of limitation.

09636801.081000

EXAMPLES

Example 1Identification of Representative Ovarian Carcinoma Protein cDNAs

5

This Example illustrates the identification of cDNA molecules encoding ovarian carcinoma proteins.

Anti-SCID mouse sera (generated against sera from SCID mice carrying late passage ovarian carcinoma) was pre-cleared of E. coli and phage antigens and used at a 1:200 dilution in a serological expression screen. The library screened was made from a SCID-derived human ovarian tumor (OV9334) using a directional RH oligo(dT) priming cDNA library construction kit and the λ Screen vector (Novagen). A bacteriophage lambda screen was employed. Approximately 400,000 pfu of the amplified OV9334 library were screened.

196 positive clones were isolated. Certain sequences that appear to be novel are provided in Figures 1A-1S and SEQ ID NOs:1 to 71. Three complete insert sequences are shown in Figures 2A-2C (SEQ ID NOs:72 to 74). Other clones having known sequences are presented in Figures 15A-15EEE (SEQ ID NOs:82 to 310). Database searches identified the following sequences that were substantially identical to the sequences presented in Figures 15A-15EEE.

These clones were further characterized using microarray technology to determine mRNA expression levels in a variety of tumor and normal tissues. Such analyses were performed using a Synteni (Palo Alto, CA) microarray, according to the manufacturer's instructions. PCR amplification products were arrayed on slides, with each product occupying a unique location in the array. mRNA was extracted from the tissue sample to be tested, reverse transcribed and fluorescent-labeled cDNA probes were generated. The microarrays were probed with the labeled cDNA probes and the slides were scanned to measure fluorescence intensity. Data was analyzed using Synteni's provided GEMtools software. The results for one clone (13695, also referred to as O8E) are shown in Figure 3.

Example 2

Identification of Ovarian Carcinoma cDNAs using Microarray Technology

5 This Example illustrates the identification of ovarian carcinoma polynucleotides by PCR subtraction and microarray analysis. Microarrays of cDNAs were analyzed for ovarian tumor-specific expression using a Synteni (Palo Alto, CA) microarray, according to the manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619, 1996 and Heller et al., *Proc. Natl. Acad. Sci.*
10 *USA* 94:2150-2155, 1997).

A PCR subtraction was performed using a tester comprising cDNA of four ovarian tumors (three of which were metastatic tumors) and a driver of cDNA from five normal tissues (adrenal gland, lung, pancreas, spleen and brain). cDNA fragments recovered from this subtraction were subjected to DNA microarray analysis where the
15 fragments were PCR amplified, adhered to chips and hybridized with fluorescently labeled probes derived from mRNAs of human ovarian tumors and a variety of normal human tissues. In this analysis, the slides were scanned and the fluorescence intensity was measured, and the data were analyzed using Synteni's GEMtools software. In general, sequences showing at least a 5-fold increase in expression in tumor cells (relative to normal
20 cells) were considered ovarian tumor antigens. The fluorescent results were analyzed and clones that displayed increased expression in ovarian tumors were further characterized by DNA sequencing and database searches to determine the novelty of the sequences.

Using such assays, an ovarian tumor antigen was identified that is a splice fusion between the human T-cell leukemia virus type I oncoprotein TAX (*see* Jin et al.,
25 *Cell* 93:81-91, 1998) and an extracellular matrix protein called osteonectin. A splice junction sequence exists at the fusion point. The sequence of this clone is presented in Figure 4 and SEQ ID NO:75. Osteonectin, unspliced and unaltered, was also identified from such assays independently.

Further clones identified by this method are referred to herein as 3f, 6b, 8e, 8h, 12c and 12h. Sequences of these clones are shown in Figures 5 to 9 and SEQ ID NOs:76 to 81. Microarray analyses were performed as described above, and are presented in Figures 10 to 14. A full length sequence encompassing clones 3f, 6b, 8e and 12h was
 5 obtained by screening an ovarian tumor (SCID-derived) cDNA library. This 2996 base pair sequence (designated O772P) is presented in SEQ ID NO:311, and the encoded 914 amino acid protein sequence is shown in SEQ ID NO:312. PSORT analysis indicates a Type 1a transmembrane protein localized to the plasma membrane.

In addition to certain of the sequences described above, this screen identified
 10 the following sequences:

Sequence	Comments
OV4vG11 (SEQ ID NO:313)	human clone 1119D9 on chromosome 20p12
OV4vB11 (SEQ ID NO:314)	human UWGC:y14c094 from chromosome 6p21
OV4vD9 (SEQ ID NO:315)	human clone 1049G16 chromosome 20q12-13.2
OV4vD5 (SEQ ID NO:316)	human KIAA0014 gene
OV4vC2 (SEQ ID NO:317)	human KIAA0084 gene
OV4vF3 (SEQ ID NO:318)	human chromosome 19 cosmid R31167
OV4VC1 (SEQ ID NO:319)	novel
OV4vH3 (SEQ ID NO:320)	novel
OV4vD2 (SEQ ID NO:321)	novel
O815P (SEQ ID NO:322)	novel
OV4vC12 (SEQ ID NO:323)	novel
OV4vA4 (SEQ ID NO:324)	novel
OV4vA3 (SEQ ID NO:325)	novel
OV4v2A5 (SEQ ID NO:326)	novel
O819P (SEQ ID NO:327)	novel
O818P (SEQ ID NO:328)	novel

Sequence	Comments
O817P (SEQ ID NO:329)	novel
O816P (SEQ ID NO:330)	novel
Ov4vC5 (SEQ ID NO:331)	novel
21721 (SEQ ID NO:332)	human lumican
21719 (SEQ ID NO:333)	human retinoic acid-binding protein II
21717 (SEQ ID NO:334)	human26S proteasome ATPase subunit
21654 (SEQ ID NO:335)	human copine I
21627 (SEQ ID NO:336)	human neuron specific gamma-2 enolase
21623 (SEQ ID NO:337)	human geranylgeranyl transferase II
21621 (SEQ ID NO:338)	human cyclin-dependent protein kinase
21616 (SEQ ID NO:339)	human prepro-megakaryocyte potentiating factor
21612 (SEQ ID NO:340)	human UPH1
21558 (SEQ ID NO:341)	human RalGDS-like 2 (RGL2)
21555 (SEQ ID NO:342)	human autoantigen P542
21548 (SEQ ID NO:343)	human actin-related protein (ARP2)
21462 (SEQ ID NO:344)	human huntingtin interacting protein
21441 (SEQ ID NO:345)	human 90K product (tumor associated antigen)
21439 (SEQ ID NO:346)	human guanine nucleotide regulator protein (tim1)
21438 (SEQ ID NO:347)	human Ku autoimmune (p70/p80) antigen
21237 (SEQ ID NO:348)	human S-laminin
21436 (SEQ ID NO:349)	human ribophorin I
21435 (SEQ ID NO:350)	human cytoplasmic chaperonin hTRiC5
21425 (SEQ ID NO:351)	humanEMX2
21423 (SEQ ID NO:352)	human p87/p89 gene
21419 (SEQ ID NO:353)	human HPBR11-7
21252 (SEQ ID NO:354)	human T1-227H
21251 (SEQ ID NO:355)	human cullin I

52

Sequence	Comments
Human arginosuccinate synth (SEQ ID NO:382)	
Human acidic ribosomal phosphoprotein (SEQ ID NO:383)	
Human colon carcinoma laminin binding pro (SEQ ID NO:384)	

This screen further identified multiple forms of the clone O772P, referred to herein as 21013, 21003 and 21008. PSORT analysis indicates that 21003 (SEQ ID NO:386; translated as SEQ ID NO:389) and 21008 (SEQ ID NO:387; translated as SEQ ID NO:390) represent Type 1a transmembrane protein forms of O772P. 21013 (SEQ ID NO:385; translated as SEQ ID NO:388) appears to be a truncated form of the protein and is predicted by PSORT analysis to be a secreted protein.

Additional sequence analysis resulted in a full length clone for O8E (2627 bp, which agrees with the message size observed by Northern analysis; SEQ ID NO:391).

10 This nucleotide sequence was obtained as follows: the original O8E sequence (OrigO8Econs) was found to overlap by 33 nucleotides with a sequence from an EST clone (IMAGE#1987589). This clone provided 1042 additional nucleotides upstream of the original O8E sequence. The link between the EST and O8E was confirmed by sequencing multiple PCR fragments generated from an ovary primary tumor library using primers to

15 the unique EST and the O8E sequence (ESTxO8EPCR). Full length status was further indicated when anchored PCR from the ovary tumor library gave several clones (AnchoredPCR cons) that all terminated upstream of the putative start methionine, but failed to yield any additional sequence information. Figure 16 presents a diagram that illustrates the location of each partial sequence within the full length O8E sequence.

20 Two protein sequences may be translated from the full length O8E. For "a" (SEQ ID NO:393) begins with a putative start methionine. A second form "b" (SEQ ID NO:392) includes 27 additional upstream residues to the 5' end of the nucleotide sequence.

Example 3

This example discloses the identification and characterization of antibody epitopes recognized by the O8E polyclonal anti-sera.

Rabbit anti-sera was raised against E. coli derived O8E recombinant protein and tested for antibody epitope recognition against 20 or 21 mer peptides that correspond to the O8E amino acid sequence. Peptides spanning amino acid regions 31 to 65, 76 to 110, 136 to 200 and 226 to 245 of the full length O8E protein were recognized by an acid eluted peak and/or a salt eluted peak from affinity purified anti-O8E sera. Thus, the corresponding amino acid sequences of the above peptides constitute the antibody epitopes recognized by affinity purified anti-O8E antibodies.

For epitope mapping, 20 or 21 mer peptides corresponding to the O8E protein were synthesized. For antibody affinity purification, rabbit anti-O8E sera was run over an O8E-sepharose column, then antibody was eluted with a salt buffer containing 0.5 M NaCl and 20 mM PO₄, followed by an acid elution step using 0.2 M Glycine, pH 2.3. Purified antibody was neutralized by the addition of 1M Tris, pH 8 and buffer exchanged into phosphate buffered saline (PBS). For enzyme linked immunosorbant assay (ELISA) analysis, O8E peptides and O8E recombinant protein were coated onto 96 well flat bottom plates at 2 µg/ml for 2 hours at room temperature (RT). Plates were then washed 5 times with PBS + 0.1 % Tween 20 and blocked with PBS + 1 % bovine serum albumin (BSA) for 1 hour. Affinity purified anti-O8E antibody, either an acid or salt eluted fraction, was then added to the wells at 1 µg/ml and incubated at RT for 1 hr. Plates were again washed, followed by the addition of donkey anti-rabbit-Ig-horseradish peroxidase (HRP) antibody for 1 hour at RT. Plates were washed, then developed by the addition of the chromagenic substrate 3, 3', 5, 5'-tetramethylbenzidine (TMB) (described by Bos *et al.*, *J. of Immunoassay* 2:187-204 (1981); available from Sigma (St. Louis, MO)). The reaction was incubated 15 minutes at RT and then stopped by the addition of 1 N H₂SO₄. Plates were read at an optical density of 450 (OD450) in an automated plate reader. The sequences of peptides corresponding to the O8E antibody epitopes are disclosed herein as SEQ ID NOs:

394-415. Antibody epitopes recognized by the O8E polyclonal anti-sera are disclosed herein in Figure 17.

Example 4

5 This example discloses IHC analysis of O8E expression in ovarian cancer tissue samples.

For immunohistochemistry studies, paraffin-embedded formalin fixed ovarian cancer tissue was sliced into 8 micron sections. Steam heat induced epitope retrieval (SHIER) in 0.1 M sodium citrate buffer (pH 6.0) was used for optimal staining
10 conditions. Sections were incubated with 10% serum/PBS for 5 minutes. Primary antibody (anti-O8E rabbit affinity purified polyclonal antibody) was added to each section for 25 min followed by a 25 min incubation with an anti-rabbit biotinylated antibody. Endogenous peroxidase activity was blocked by three 1.5 min incubations with hydrogen peroxidase. The avidin biotin complex/horse radish peroxidase system was used along
15 with DAB chromogen to visualize antigen expression. Slides were counterstained with hematoxylin. One (papillary serous carcinoma) of six ovarian cancer tissue sections displayed O8E immunoreactivity. O8E expression was localized to the plasma membrane.

Six ovarian cancer tissues were analyzed with the anti-O8E rabbit polyclonal antibody. One (papillary serous carcinoma) of six ovarian cancer tissue samples
20 stained positive for O8E expression. O8E expression was localized to the surface membrane.

Example 5

25 This example discloses O8E peptides that are predicted to bind HLA-A2 and to be immunogenic for CD8 T cell responses in humans.

Potential HLA-A2 binding peptides of O8E were predicted by using the full-length open-reading frame (ORF) from O8E and running it through "Episeek," a program used to predict MHC binding peptides. The program used is based on the algorithm published by Parker, K.C. *et al.*, *J. Immunol.* 152(1):163-175 (1994) (incorporated by

reference herein in its entirety). 10-mer and 9-mer peptides predicted to bind HLA-0201 are disclosed herein as SEQ ID NOs: 416-435 and SEQ ID NOs: 436-455, respectively.

Example 6

5 This example discloses O8E cell surface expression measured by fluorescence activated cell sorting.

For FACS analysis, cells were washed with ice cold staining buffer (PBS/1% BSA/azide). Next, the cells were incubated for 30 minutes on ice with 10 micrograms/ml of affinity purified rabbit anti-B305D polyclonal antibody. The cells were
10 washed 3 times with staining buffer and then incubated with a 1:100 dilution of a goat anti-rabbit Ig (H+L)-FITC reagent (Southern Biotechnology) for 30 minutes on ice. Following 3 washes, the cells were resuspended in staining buffer containing prodium iodide, a vital stain that allows for identification of permeable cells, and analyzed by FACS. O8E surface expression was confirmed on SKBR3 breast cancer cells and HEK293
15 cells that stably overexpress the cDNA for O8E. Neither MB415 cells nor HEK293 cells stably transfected with a control irrelevant plasmid DNA showed surface expression of O8E (Figures 18 and 19).

Example 7

This example further evaluates the expression and surface localization of
20 O8E.

For expression and purification of antigen used for immunization, O8E expressed in an E. coli recombinant expression system was grown overnight in LB Broth with the appropriate antibiotics at 37°C in a shaking incubator. The next morning, 10 ml of the overnight culture was added to 500 ml of 2x YT plus appropriate antibiotics in a 2L-
25 baffled Erlenmeyer flask. When the Optical Density (at 560 nanometers) of the culture reached 0.4-0.6 the cells were induced with IPTG (1 mM). 4 hours after induction with IPTG the cells were harvested by centrifugation. The cells were then washed with phosphate buffered saline and centrifuged again. The supernatant was discarded and the cells were either frozen for future use or immediately processed. Twenty milliliters of lysis

buffer was added to the cell pellets and vortexed. To break open the E. coli cells, this mixture was then run through the French Press at a pressure of 16,000 psi. The cells were then centrifuged again and the supernatant and pellet were checked by SDS-PAGE for the partitioning of the recombinant protein. For protein that localized to the cell pellet, the

5 pellet was resuspended in 10 mM Tris pH 8.0 , 1% CHAPS and the inclusion body pellet was washed and centrifuged again. This procedure was repeated twice more. The washed inclusion body pellet was solubilized with either 8 M urea or 6 M guanidine HCl containing 10 mM Tris pH 8.0 plus 10 mM imidazole. The solubilized protein was added to 5 ml of nickel-chelate resin (Qiagen) and incubated for 45 min to 1 hour at room

10 temperature with continuous agitation. After incubation, the resin and protein mixture were poured through a disposable column and the flow through was collected. The column was then washed with 10-20 column volumes of the solubilization buffer. The antigen was then eluted from the column using 8M urea, 10 mM tris pH 8.0 and 300 mM imidazole and collected in 3 ml fractions. A SDS-PAGE gel was run to determine which fractions to pool

15 for further purification. As a final purification step, a strong anion exchange resin such as Hi-Prep Q (Biorad) was equilibrated with the appropriate buffer and the pooled fractions from above were loaded onto the column. Each antigen was eluted off of the column with an increasing salt gradient. Fractions were collected as the column was run and another SDS-PAGE gel was run to determine which fractions from the column to pool. The pooled

20 fractions were dialyzed against 10 mM Tris pH 8.0. This material was then evaluated for acceptable purity as determined by SDS-PAGE or HPLC, concentration as determined by Lowry assay or Amino Acid Analysis, identity as determined by amino terminal protein sequence, and endotoxin level as determined by the Limulus (LAL) assay. The proteins were then vialled after filtration through a 0.22 micron filter and the antigens were frozen

25 until needed for immunization.

For generation of polyclonal anti-sera, 400 micrograms of each prostate antigen was combined with 100 micrograms of muramyl dipeptide (MDP). Equal volume of Incomplete Freund's Adjuvant (IFA) was added and then mixed. Every four weeks animals were boosted with 100 micrograms of antigen mixed with an equal volume of IFA. Seven

days following each boost the animal was bled. Sera was generated by incubating the blood at 4°C for 12-24 hours followed by centrifugation.

For characterization of polyclonal antisera, 96 well plates were coated with antigen by incubating with 50 microliters (typically 1 microgram) at 4 C for 20 hrs. 250 microliters of BSA blocking buffer was added to the wells and incubated at RT for 2 hrs. Plates were washed 6 times with PBS/0.01% tween. Anti-O8E rabbit sera or affinity purified anti-O8e antibody was diluted in PBS. Fifty microliters of diluted antibody was added to each well and incubated at RT for 30 min. Plates were washed as described above before 50 microliters of goat anti-rabbit horse radish peroxidase (HRP) at a 1:10000 dilution was added and incubated at RT for 30 min. Plates were washed as described above and 100 microliters of TMB microwell Peroxidase Substrate was added to each well. Following a 15 minute incubation in the dark at room temperature the colorimetric reaction was stopped with 100 microliters of 1N H₂SO₄ and read immediately at 450 nm. All polyclonal antibodies showed immunoreactivity to the O8E antigen.

For recombinant expression in mammalian HEK293 cells, full length O8E cDNA was subcloned into the mammalian expression vectors pcDNA3.1+ and pCEP4 (Invitrogen) which were modified to contain His and FLAG epitope tags, respectively. These constructs were transfected into HEK293 cells (ATCC) using Fugene 6 reagent (Roche). Briefly, HEK293 cells were plated at a density of 100,000 cells/ml in DMEM (Gibco) containing 10% FBS (Hyclone) and grown overnight. The following day, 2 ul of Fugene6 was added to 100 ul of DMEM containing no FBS and incubated for 15 minutes at room temperature. The Fugene6/DMEM mixture was then added to 1ug of O8E/pCEP4 or O8E/pcDNA3.1 plasmid DNA and incubated for 15 minutes at room temperature. The Fugene/DNA mix was then added to the HEK293 cells and incubated for 48-72 hrs at 37°C with 7% CO₂. Cells were rinsed with PBS then collected and pelleted by centrifugation. For Western blot analysis, whole cell lysates were generated by incubating the cells in Triton-X100 containing lysis buffer for 30 minutes on ice. Lysates were then cleared by centrifugation at 10,000rpm for 5 minutes at 4 C. Samples were diluted with SDS-PAGE loading buffer containing beta-mercaptoethanol, then boiled for 10 minutes

prior to loading the SDS-PAGE gel. Protein was transferred to nitrocellulose and probed using anti-O8E rabbit polyclonal sera #2333L at a dilution of 1:750. The blot was revealed with a goat anti-rabbit Ig coupled to HRP followed by incubation in ECL substrate.

For FACS analysis, cells were washed further with ice cold staining buffer (PBS+1%BSA+Azide). Next, the cells were incubated for 30 minutes on ice with 10ug/ml of Protein A purified anti-O8E polyclonal sera. The cells were washed 3 times with staining buffer and then incubated with a 1:100 dilution of a goat anti-rabbit Ig(H+L)-FITC reagent (Southern Biotechnology) for 30 minutes on ice. Following 3 washes, the cells were resuspended in staining buffer containing Propidium Iodide (PI), a vital stain that allows for the identification of permeable cells, and analyzed by FACS.

From these experiments, the results of which are illustrated in Figures 20-21, O8E expression was detected on the surface of transfected HEK293 cells and SKBR3 cells by FACS analysis using rabbit anti-O8E sera. Expression was also detected in transfected HEK293 cell lysates by Western blot analysis (not shown).

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

SUMMARY OF SEQUENCE LISTING

SEQ ID NOs:1-71 are ovarian carcinoma antigen polynucleotides shown in Figures 1A-1S.

SEQ ID NOs:72-74 are ovarian carcinoma antigen polynucleotides shown in Figures 2A-2C.

SEQ ID NO:75 is the ovarian carcinoma polynucleotide 3g (Figure 4).

SEQ ID NO:76 is the ovarian carcinoma polynucleotide 3f (Figure 5).

SEQ ID NO:77 is the ovarian carcinoma polynucleotide 6b (Figure 6).

SEQ ID NO:78 is the ovarian carcinoma polynucleotide 8e (Figure 7A).

SEQ ID NO:79 is the ovarian carcinoma polynucleotide 8h (Figure 7B).

SEQ ID NO:80 is the ovarian carcinoma polynucleotide 12e (Figure 8).

SEQ ID NO:81 is the ovarian carcinoma polynucleotide 12h (Figure 9).

SEQ ID NOs:82-310 are ovarian carcinoma antigen polynucleotides shown
5 in Figures 15A-15EEE.

SEQ ID NO:311 is a full length sequence of ovarian carcinoma
polynucleotide O772P.

SEQ ID NO:312 is the O772P amino acid sequence.

SEQ ID NOs:313-384 are ovarian carcinoma antigen polynucleotides.

10 SEQ ID NOs:385-390 present sequences of O772P forms.

SEQ ID NO:391 is a full length sequence of ovarian carcinoma
polynucleotide O8E.

SEQ ID NOs:392-393 are protein sequences encoded by O8E.

09536801-081000